

Proof of the De Concini-Kac-Procesi conjecture

Kobi Kremnizer

February 2, 2008

Abstract

In this paper we prove a conjecture by De Concini, Kac and Procesi [CP] (Corollary 5.3):

Corollary. *The dimension of any $M \in U_q - \text{mod}^\chi$ is divisible by $l^{\text{codim}_{\mathcal{B}} \mathcal{B}_\chi}$.*

1 Introduction

In this paper we prove a conjecture by De Concini, Kac and Procesi [CP] (Corollary 5.3):

Corollary. *The dimension of any $M \in U_q - \text{mod}^\chi$ is divisible by $l^{\text{codim}_{\mathcal{B}} \mathcal{B}_\chi}$.*

The conjecture follows from the following theorem (Theorem 5.1):

Theorem. *Fix χ a nilpotent element in the big cell and a regular $\lambda \in P$. For any module $M \in U_q - \text{mod}^{\lambda, \chi}$ there exists a polynomial $\mathbf{d}_M \in \frac{1}{R}\mathbb{Z}[P^*]$ of degree $\leq \dim(\mathcal{B}_\chi)$, such that for any $\mu \in P$ in the closure of the alcove of λ , we have*

$$\dim(T_\lambda^\mu(M)) = \mathbf{d}_M(\mu) \quad (1.1)$$

$\mathbf{d}_M(\mu) = l^{\dim(\mathcal{B})} \mathbf{d}_M^0(\frac{\mu+\rho}{l})$ for another polynomial $\mathbf{d}_M^0 \in \frac{1}{R}\mathbb{Z}[P^]$, such that $\mathbf{d}_M^0(\mu) \in \mathbb{Z}$ for $\mu \in P$.*

The proof of this theorem follows quite easily from the results of [BK2] and some usual arguments concerning the Euler characteristic.

It must be emphasized that [BMR] prove a similar result in positive characteristic (the Kac-Weisfeiler conjecture first proved by Premet) using geometric methods. This allowed us to adapt their proof to the quantum case by using the geometric construction of modules for the quantum group from [BK2]. Thus this paper is another indication of the close relationship between positive characteristic representations and root of unity representations. This relationship will be investigated more in the future.

We would like to thank Dmitriy Rumynin that suggested we should prove this conjecture.

The author was supported in part by NSF grant DMS-0602007.

2 Quantum groups

2.1 Conventions

Let $\mathbb{C}$ be the field of complex numbers and fix $q \in \mathbb{C}^*$.

We always assume that if q is a root of unity it is primitive of odd order and in case G has a component of type G_2 the order is also prime to 3. Let $\mathcal{A}$ be the local ring $\mathbb{Z}[\nu]_{\mathfrak{m}}$, where $\mathfrak{m}$ is the maximal ideal in $\mathbb{Z}[\nu]$ generated by $\nu - 1$ and a fixed odd prime p .

2.2 Root data

Let $\mathfrak{g}$ be a semi-simple Lie algebra and let $\mathfrak{h} \subset \mathfrak{b}$ be a Cartan subalgebra contained in a Borel subalgebra of $\mathfrak{g}$. Let R be the root system, $\Delta \subset R_+ \subset R$ a basis and the positive roots. Let $P \subset \mathfrak{h}^*$ be the weight lattice and P_+ the positive weights; the i 'th fundamental weight is denoted by ω_i and ρ denotes the half sum of the positive roots. Let $Q \subset P$ be the root lattice and $Q_+ \subset Q$ those elements which have non-negative coefficients with respect to the basis of simple roots. Let $\mathcal{W}$ be the Weyl group of $\mathfrak{g}$. We let $\langle, \rangle$ denote a $\mathcal{W}$ -invariant bilinear form on $\mathfrak{h}^*$ normalized by $\langle \gamma, \gamma \rangle = 2$ for each short root γ .

Let $T_P = \text{Hom}_{\text{groups}}(P, k^*)$ be the character group of P with values in k (we use additive notation for this group). If $\mu \in P$, then $\langle \mu, P \rangle \subset \mathbb{Z}$ and hence we can define $q^\mu \in T_P$ by the formula $q^\mu(\gamma) = q^{\langle \mu, \gamma \rangle}$, for $\gamma \in P$. If

$\mu \in P, \lambda \in T_P$ we write $\mu + \lambda = q^\mu + \lambda$. Note that the Weyl group naturally acts on T_P .

2.3 Quantized enveloping algebra U_q and quantized algebra of functions $\mathcal{O}_q$.

Let U_q be the simply connected quantized enveloping algebra of $\mathfrak{g}$ over $\mathbb{C}$. Recall that U_q has algebra generators $E_\alpha, F_\alpha, K_\mu, \alpha, \beta$ are simple roots, $\mu \in P$ subject to the relations

$$K_\lambda K_\mu = K_{\lambda+\mu}, \quad K_0 = 1, \quad (2.1)$$

$$K_\mu E_\alpha K_{-\mu} = q^{<\mu, \alpha>} E_\alpha, \quad K_\mu F_\alpha K_{-\mu} = q^{-<\mu, \alpha>} F_\alpha, \quad (2.2)$$

$$[E_\alpha, F_\beta] = \delta_{\alpha, \beta} \frac{K_\alpha - K_{-\alpha}}{q_\alpha - q_\alpha^{-1}} \quad (2.3)$$

and certain Serre-relations that we do not recall here. Here

$q_\alpha = q^{d_\alpha}$, $d_\alpha = <\alpha, \alpha> / 2$. (We have assumed that $q_\alpha^2 \neq 1$.)

Let G be the simply connected algebraic group with Lie algebra $\mathfrak{g}$, B be a Borel subgroup of G and $N \subset B$ its unipotent radical. Let $\mathfrak{b} = \text{Lie } B$ and $\mathfrak{n} = \text{Lie } N$ and denote by $U_q(\mathfrak{b})$ and $U_q(\mathfrak{n})$ the corresponding subalgebras of U_q . Then $U_q(\mathfrak{b})$ is a Hopf algebra, while $U_q(\mathfrak{n})$ is only an algebra. Let $\mathcal{O}_q = \mathcal{O}_q(G)$ be the algebra of matrix coefficients of finite dimensional type-1 representations of U_q . There is a natural pairing $(,) : U_q \otimes \mathcal{O}_q \rightarrow \mathbb{C}$. This gives a U_q -bimodule structure on $\mathcal{O}_q$ as follows

$$ua = a_1(u, a_2), \quad au = (u, a_1)a_2, \quad u \in U_q, a \in \mathcal{O}_q \quad (2.4)$$

Then $\mathcal{O}_q$ is the (restricted) dual of U_q with respect to this pairing. We let $\mathcal{O}_q(B)$ and $\mathcal{O}_q(N)$ be the quotient algebras of $\mathcal{O}_q$ corresponding to the subalgebras $U_q(\mathfrak{b})$ and $U_q(\mathfrak{n})$ of U_q , respectively, by means of this duality. Then $\mathcal{O}_q(B)$ is a Hopf algebra and $\mathcal{O}_q(N)$ is only an algebra.

There is a braid group action on U_q . For each $w \in \mathcal{W}$, we get an automorphism T_w of U_q .

2.4 Integral versions of U_q .

Let U_q^{res} be the Lusztig's integral form of U_q , the $\mathcal{A}$ -algebra in U_q generated by divided powers $E_\alpha^{(n)} = E_\alpha^n / [n]_{d_\alpha}!$, $F_\alpha^{(n)} = F_\alpha^n / [n]_{d_\alpha}!$, α a simple root,

$n \geq 1$ (where $[m]_d = \frac{\prod_{s=1}^m (q^{d \cdot s} - q^{-d \cdot s})}{q^d - q^{-d}}$) and the K_μ 's, $\mu \in P$. There is also the De Consini-Kac integral form $U_{\mathcal{A}}$, which is generated over $\mathcal{A}$ by the E_α, F_α and K_μ 's. The subalgebra $U_{\mathcal{A}}$ is preserved by the adjoint action of $U_{\mathcal{A}}^{res}$: $ad_{U_{\mathcal{A}}^{res}}(U_{\mathcal{A}}) \subset U_{\mathcal{A}}$. The operators T_w from section 2.1.2 preserves the integral versions.

$\mathcal{O}_{\mathcal{A}}$ is defined to be the dual of $U_{\mathcal{A}}^{res}$. This is an $\mathcal{A}$ -sub Hopf algebra of $\mathcal{O}_q$.

2.5 Finite part of U_q .

The algebra U_q acts on itself by the adjoint action $ad : U_q \rightarrow U_q$ where $ad(u)(v) = u_1 v S(u_2)$. Let U_q^{fin} be the finite part of U_q with respect to this action:

$$U_q^{fin} = \{v \in U_q; \dim ad(U_q)(v) < \infty\}.$$

This is a subalgebra. (See [JL].)

We can also give an integral version of the finite part as the finite part of the action of $U_{\mathcal{A}}^{res}$ on $U_{\mathcal{A}}$. Thus by specializing we get a subalgebra of U_q for every q . Of course, when specialized to generic q this coincides with the previous definition.

2.6 Specializations and Frobenius maps.

For any ring map $\phi : \mathcal{A} \rightarrow R$ we put $U_R = U_{\mathcal{A}} \otimes_{\mathcal{A}} R$ and $U_R^{res} = U_{\mathcal{A}}^{res} \otimes_{\mathcal{A}} R$. If $R = \mathbb{C}$ and $\phi(\nu) = q$, there are three different cases: q is a root of unity, $q = 1$ and q is generic. Then $U_R = U_q$.

There is also the ring map $\mathcal{A} \rightarrow \mathbb{F}_p$, sending $\nu \rightarrow 1$. Then $U_{\mathbb{F}_p}/(K - 1) = U(\mathfrak{g}_p)$, the enveloping algebra of the Lie algebra $\mathfrak{g}_p$ in characteristic p .

For any $U_{\mathcal{A}}$ -module (resp. $U_{\mathcal{A}}^{res}$ -module) $M_{\mathcal{A}}$ we put $M_R = M_{\mathcal{A}} \otimes_{\mathcal{A}} R$. This is an U_R -module (resp. U_R^{res} -module). When $R = \mathbb{C}$ we simply write $M = M_{\mathbb{C}}$.

When q is a root of unity, we have the Frobenius map: $U_q^{res} \rightarrow U(\mathfrak{g})$. Its algebra kernel is denoted by $\mathfrak{u}_q$. We also have the Frobenius map $U_q^{res}(\mathfrak{b}) \rightarrow U(\mathfrak{b})$, with algebra kernel b_q . These maps induces dual maps $\mathcal{O} = \mathcal{O}(G) \hookrightarrow \mathcal{O}_q$ and $\mathcal{O}(B) \hookrightarrow \mathcal{O}_q(B)$.

For each q there exists a map $U_q \rightarrow U_q^{res}$ whose image is $\mathfrak{u}_q$ and whose algebra kernel is $\mathcal{Z}^{(l)}$ (see section 2.1.7 below for the definition of $\mathcal{Z}^{(l)}$).

2.7 Verma modules.

For each $\lambda \in T_P$ there is the one dimensional $U_q(\mathfrak{b})$ -module $\mathbb{C}_\lambda$ which is given by extending λ to act by zero on the E_α 's. The Verma-module M_λ is the U_q -module induced from $\mathbb{C}_\lambda$. If $\mu \in P$ we write $M_\mu = M_{q^\mu}$. A point important for us is that M_λ carries an $U_q^{res}(\mathfrak{b})$ -module structure defined as follows: $U_q^{res}(\mathfrak{b})$ acts on U_q by restricting the adjoint action of U_q^{res} on U_q . This induces a $U_q(\mathfrak{b})$ -action on the quotient M_λ of U_q . Since this action is locally finite it corresponds to an $\mathcal{O}_q(B)$ -comodule action on M_λ . **Note!!** As a $U_q^{res}(\mathfrak{b})$ -module M_λ has *trivial highest weight*. (In case q is generic, $U_q^{res}(\mathfrak{b}) = U_q(\mathfrak{b})$ and then the $U_q^{res}(\mathfrak{b})$ action on M_λ described above is the same as the $U(\mathfrak{b})$ -action on $M_\lambda \otimes \mathbb{C}_{-\lambda}$.)

Verma module M_λ has an integral version $M_{\lambda, A}$.

2.8 Centers of U_q and definition of $\tilde{U}_q$.

Let $\mathcal{Z}$ denote the center of U_q . When q is a p 'th root of unity $\mathcal{Z}$ contains the Harish-Chandra center $\mathcal{Z}^{HC}$ and the l 'th center $\mathcal{Z}^{(l)}$ which is generated by the E_α^l , F_α^l , K_μ^l and K_μ^{-l} 's. In fact, $\mathcal{Z} = \mathcal{Z}^{(l)} \otimes_{\mathcal{Z}^{(l)} \cap \mathcal{Z}^{HC}} \mathcal{Z}^{HC}$. There is the Harish-Chandra homomorphism $\mathcal{Z}^{HC} \rightarrow \mathcal{O}(T_P)$ that maps isomorphically to the W -invariant even part of $\mathcal{O}(T_P)$. We define $\tilde{U}_q = U_q \otimes_{\mathcal{Z}^{HC}} \mathcal{O}(T_P)$.

2.9 Some conventions.

We shall frequently refer to a right (resp. left) $\mathcal{O}_q$ -comodule as a left (resp. right) G_q -module, etc. If we have two right $\mathcal{O}_q$ -comodules V and W , then $V \otimes W$ carries the structure of a right $\mathcal{O}_q$ -comodule via the formula

$$\delta(v \otimes w) = v_1 \otimes w_1 \otimes v_2 w_2$$

We shall refer to this action as the *tensor* or *diagonal* action. A similar formula exist for left comodules.

3 Quantum flag variety

Here we recall the definition and basic properties of the quantum flag variety from [BK1].

3.0.1 Category $\mathcal{M}_{B_q}(G_q)$.

The composition

$$\mathcal{O}_q \rightarrow \mathcal{O}_q \otimes \mathcal{O}_q \rightarrow \mathcal{O}_q \otimes \mathcal{O}_q(B) \quad (3.1)$$

defines a right $\mathcal{O}_q(B)$ -comodule structure on $\mathcal{O}_q$. A B_q -equivariant sheaves on G_q is a triple (F, α, β) where F is a vector space, $\alpha : \mathcal{O}_q \otimes F \rightarrow F$ a left $\mathcal{O}_q$ -module action and $\beta : F \rightarrow F \otimes \mathcal{O}_q(B)$ a right $\mathcal{O}_q(B)$ -comodule action such that α is a right comodule map, where we consider the tensor comodule structure on $\mathcal{O}_q(G) \otimes F$.

Definition 3.1. We denote $\mathcal{M}_{B_q}(G_q)$ to be the category of B_q -equivariant sheaves on G_q . Morphisms in $\mathcal{M}_{B_q}(G_q)$ are those compatible with all structures.

If $q = 1$, the category $\mathcal{M}_B(G)$ is equivalent to the category $\mathcal{M}(G/B)$ of quasi-coherent sheaves on $\mathcal{B} = G/B$.

Definition 3.2. We define the induction functor $\text{Ind} : B_q\text{-mod} \rightarrow \mathcal{M}_{B_q}(G_q)$, $\text{Ind } V = \mathcal{O}_q \otimes V$ with the tensor B_q -action and the $\mathcal{O}_q$ -action on the first factor. For $\lambda \in P$ we define a line bundle $\mathcal{O}_q(\lambda) = \text{Ind } \mathbb{C}_{-\lambda}$.

Definition 3.3. The global section functor $\Gamma : \mathcal{M}_{B_q}(G_q) \rightarrow \mathbb{C}\text{-mod}$ is defined by

$$\Gamma(M) = \text{Hom}_{\mathcal{M}_{B_q}(G_q)}(\mathcal{O}_q, M) = \{m \in M; \Delta_B(m) = m \otimes 1\}.$$

This is the set of B_q -invariants in M .

The category $\mathcal{M}_{B_q}(G_q)$ has enough injectives, so derived functors $R\Gamma$ are well-defined. We showed that $R^i\Gamma(\text{Ind } V) = H^i(G_q/B_q, V)$, where $H^i(G_q/B_q, \cdot)$ is the i 'th derived functor of the functor $V \rightarrow \Gamma(\text{Ind } V)$ from $B_q\text{-mod}$ to $\mathbb{C}\text{-mod}$.

We proved a quantum version of Serre's basic theorem on projective schemes: Each $M \in \mathcal{M}_{B_q}(G_q)$ is a quotient of a direct sum of $\mathcal{O}_q(\lambda)$'s and each surjection $M \twoheadrightarrow M'$ of noetherian objects in $\mathcal{M}_{B_q}(G_q)$ induces a surjection $\Gamma(M(\lambda)) \twoheadrightarrow \Gamma(M'(\lambda))$ for $\lambda \gg 0$.

Here the notation $\lambda \gg 0$ means that $\langle \lambda, \alpha^\vee \rangle$ is a sufficiently large integer for each simple root α and $M(\lambda) = M \otimes \mathbb{C}_{-\lambda}$ is the λ -twist of M .

Let $V \in G_q\text{-mod}$. Denote by $V|_{B_q}$ the restriction of V to B_q and by V^{triv} the trivial B_q -module whose underlying space is V . We showed that $\text{Ind } V|_{B_q}$ and $\text{Ind } V^{triv}$ are isomorphic in $\mathcal{M}_{B_q}(G_q)$. In particular

$$\Gamma(\text{Ind } V|_{B_q}) = V|_{B_q} \otimes \Gamma(\mathcal{O}_q) = V|_{B_q}, \text{ for } V \in G_q\text{-mod} \quad (3.2)$$

3.0.2 $\mathcal{M}_{B_q}(G_q)$ at a root of unity

In case q is a root of unity we have the following Frobenius morphism:

$$Fr_* : \mathcal{M}_{B_q}(G_q) \rightarrow \mathcal{M}(G/B) \quad (3.3)$$

defined as

$$N \mapsto N^{b_q} \quad (3.4)$$

Using the description of $\mathcal{M}_{B_q}(G_q)$ as $Proj(A_q)$ where $A_q = \bigoplus V_{q,\lambda}$ [BK1] and similarly $\mathcal{M}(G/B) = Proj(A)$ where $A = \bigoplus V_\lambda$, we see that Fr_* is induced from the quantum Frobenius map $A \hookrightarrow A_q$. It follows that:

Proposition 3.4. *Fr_* is exact and faithful.*

This functor has a left adjoint which we denote by Fr^* .

We will need the following:

Lemma 3.5. $Fr^*(\mathcal{F}(\lambda)) \simeq Fr^*(\mathcal{F})(l\lambda)$

Lemma 3.6. $Fr_*(Fr^*(\mathcal{F})(\lambda)) \simeq \mathcal{F} \otimes Fr_*(\mathcal{O}_{q,\lambda})$

Lemma 3.7. $Fr_*(\mathcal{O}_q(-\rho)) \simeq \mathcal{O}(-\rho)^{l^{dim(\mathcal{B})}}$

4 The ring $\mathcal{D}_{G_q}$, the category of $\mathcal{D}_q^\lambda$ -modules and translation functors

4.1 $\mathcal{D}_q^\lambda$ -modules

We need the following important

Remark 4.1. All objects described in the preceding chapters are defined over $\mathcal{A}$. For any specialization $\mathcal{A} \rightarrow R$ and any object Obj we denote by Obj_R its R -form. For the functors we don't use any subscripts; so, for instance, there is the functor $\text{Ind} : B_{q,R} - \text{mod} \rightarrow \mathcal{M}_{B_q}(G_q)_R$.

Recall the U_q -bimodule structure on $\mathcal{O}_q$ given by 2.4. Now, as we have two versions of the quantum group we pick the following definition of the ring of differential operators on the group (the *crystalline* version).

Definition 4.2. We define the ring of quantum differential operators on G_q to be the smash product algebra $\mathcal{D}_{G_q} := \mathcal{O}_q \star U_q$. So $\mathcal{D}_{G_q} = \mathcal{O}_q \otimes U_q$ as a vector space and multiplication is given by

$$a \otimes u \cdot b \otimes v = au_1(b) \otimes u_2v. \quad (4.1)$$

We consider now the ring $\mathcal{D}_{G_q}$ as a left U_q^{res} -module, via the left U_q^{res} -action on $\mathcal{O}_q$ in 2.4 and the left adjoint action of U_q^{res} on U_q ; this way $\mathcal{D}_{G_q}$ becomes a module algebra for U_q^{res} : In the following we will use the restriction of this action to $U_q^{res}(\mathfrak{b}) \subset U_q^{res}$. As U_q is not locally finite with respect to the adjoint action, this $U_q^{res}(\mathfrak{b})$ -action doesn't integrate to a B_q -action. Thus $\mathcal{D}_{G_q}$ is not an object of $\mathcal{M}_{B_q}(G_q)$; however, $\mathcal{D}_{G_q}$ has a subalgebra $\mathcal{D}_{G_q}^{fin} = \mathcal{O}_q \star U_q^{fin}$ which belongs to $\mathcal{M}_{B_q}(G_q)$. This fact will be used below.

Definition 4.3. Let $\lambda \in T_P$. A (B_q, λ) -equivariant $\mathcal{D}_{G_q}$ -module is a triple (M, α, β) , where M is a $\mathbb{C}$ -module, $\alpha : \mathcal{D}_{G_q} \otimes M \rightarrow M$ a left $\mathcal{D}_{G_q}$ -action and $\beta^{res} : M \rightarrow M \otimes \mathcal{O}_q(B)$ a right $\mathcal{O}_q(B)$ -coaction. The latter action induces an $U_q^{res}(\mathfrak{b})$ -action on M again denoted by β^{res} . We have the natural map $U_q(\mathfrak{b}) \rightarrow U_q^{res}(\mathfrak{b})$ which together with β^{res} gives an action β of $U_q(\mathfrak{b})$ on M . We require

- i) The $U_q(\mathfrak{b})$ -actions on $M \otimes \mathbb{C}_\lambda$ given by $\beta \otimes \lambda$ and by $(\alpha|_{U_q(\mathfrak{b})}) \otimes \text{Id}$ coincide.
- ii) The map α is $U_q^{res}(\mathfrak{b})$ -linear with respect to the β -action on M and the action on $\mathcal{D}_{G_q}$.

These objects form a category denoted $\mathcal{D}_{B_q}^\lambda(G_q)$. There is the forgetful functor $\mathcal{D}_{B_q}^\lambda(G_q) \rightarrow \mathcal{M}_{B_q}(G_q)$. Morphisms in $\mathcal{D}_{B_q}^\lambda(G_q)$ are morphisms in $\mathcal{M}_{B_q}(G_q)$ that are $\mathcal{D}_{G_q}$ -linear.

We defined $\mathcal{D}_q^\lambda$ as the maximal quotient of $\mathcal{D}_{G_q}$ which is an object of $\mathcal{D}_{B_q}^\lambda(G_q)$ and showed that

$$\mathcal{D}_q^\lambda = \text{Ind } M_\lambda \quad (4.2)$$

as an object in $\mathcal{D}_{B_q}^\lambda(G_q)$. (See section 2.7 for the B_q -action $= U_q^{res}(\mathfrak{b})$ -action on M_λ). The global section functor $\Gamma : \mathcal{D}_{B_q}^\lambda(G_q) \rightarrow \mathcal{M}$ is the functor of taking B_q invariants (with respect to the action β); we have $\Gamma = \text{Hom}_{\mathcal{D}_{B_q}^\lambda(G_q)}(\mathcal{D}_q^\lambda, \cdot)$.

Hence, in particular $\Gamma(\mathcal{D}_q^\lambda) = \text{End}_{\mathcal{D}_{B_q}^\lambda(G_q)}(\mathcal{D}_q^\lambda)$ (which explains the ring structure on $\Gamma(\mathcal{D}_q^\lambda)$).

Let q be a l -th root of unity. In [BK2] the following were proved:

Proposition 4.4. *We have i) $U_q^\lambda \cong R\Gamma(\mathcal{D}_q^\lambda)$ and ii) $\tilde{U}_q \cong R\Gamma(\tilde{\mathcal{D}}_q)$ (if l is a prime $p >$ Coxeter number of G .)*

Proposition 4.5. $\mathcal{D}_q^\lambda$ is an Azumaya algebra over a dense subset of T^*X^λ .

Proposition 4.6. *The action map $U_q^{fin} \otimes_Z \mathcal{O}_{\tilde{T}^*X}^{twist} \rightarrow \tilde{\mathcal{D}}_q$ induces an isomorphism $U_q^{fin} \otimes_Z \mathcal{O}_{\tilde{T}^*X}^{twist, unram} \cong \tilde{\mathcal{D}}_q|_{\tilde{T}^*X}^{twist, unram}$.*

Theorem 4.7. $R\Gamma : D^b(\mathcal{D}_{B_q}^\lambda(G_q)) \rightarrow D^b(U_q^\lambda - \text{mod})$ is an equivalence of categories.

4.2 Translation Functors

For $\lambda, \mu \in P$ define the translation functor $T_\lambda^\mu : U_q^\lambda - \text{mod} \rightarrow U_q^\mu - \text{mod}$ $T_\lambda^\mu(M) = [V_{\mu-\lambda} \otimes M]_\mu$. Here $V_{\mu-\lambda}$ is the standard G_q -module with an extremal weight $\mu - \lambda$. This functor is also defined on the larger categories $U_q - \text{mod}^\lambda$ of modules with a generalized central character.

We want to define translation functors on $\mathcal{D}$ -modules as well. We first define a category $\tilde{\mathcal{D}}_{B_q}(G_q)$ of $\mathcal{D}_{G_q}$ -modules that contains all $\mathcal{D}_q^\lambda$, $\lambda \in T_P$, a "torsor".

Definition 4.8. An object of $\tilde{\mathcal{D}}_{B_q}(G_q)$ is a triple (M, α, β) , where $\alpha : \mathcal{D}_{G_q} \otimes M \rightarrow M$ a left $\mathcal{D}_{G_q}$ -action and $\beta^{res} : M \rightarrow M \otimes \mathcal{O}_q(B)$ a right $\mathcal{O}_q(B)$ -coaction.

- i) The $U_q(\mathfrak{n}_+)$ -actions on M given by $\beta|_{U_q(\mathfrak{n}_+)}$ and by $(\alpha|_{U_q(\mathfrak{n}_+)})$ coincide.
- ii) The map α is $U_q(\mathfrak{b})$ -linear with respect to the β -action on M and the action on $\mathcal{D}_{G_q}$.

The functor of B_q -invariants (global sections) will be denoted $\tilde{\Gamma}$. Put $\tilde{\mathcal{M}} = U_q(\mathfrak{g}) / \sum_{\alpha \in R_+} U_q(\mathfrak{g}) \cdot E_\alpha$ (a "universal" Verma module) and define

$$\tilde{\mathcal{D}}_q = \mathcal{O}_q \otimes \tilde{\mathcal{M}} \quad (4.3)$$

$\tilde{\mathcal{D}}_q$ inherits an $U_q^{res}(\mathfrak{b})$ -module structure from $\mathcal{D}_{G_q}$, so $\tilde{\mathcal{D}}_q$ is an object in $\tilde{\mathcal{D}}_{B_q}(G_q)$.

We define a category $\tilde{\mathcal{D}}_{B_q}(G_q)_\lambda$ by also requiring that a power of the ideal defined by λ kills the module. We will denote the functor of B_q -invariants (global sections) by $\tilde{\Gamma}_\lambda$.

Note that we have a pair of exact functors:

$$i_\lambda : \widetilde{\mathcal{D}}_{B_q}(G_q)_\lambda \rightleftarrows \widetilde{\mathcal{D}}_{B_q}(G_q) : \llbracket_\lambda \quad (4.4)$$

where i_λ is the natural inclusion and $\llbracket_\lambda$ is the projection.

For any B_q -module V and any $\mathcal{D}_{B_q}^\lambda(G_q)$ -module M we can define a $\widetilde{\mathcal{D}}_{B_q}(G_q)$ -module $M \otimes V$ where we just twist the B_q action and let $\mathcal{D}_{G_q}$ act on the first factor. We have the following equalities:

$$T_\lambda^\mu(\widetilde{R}\Gamma_\lambda(M)) = [\widetilde{R}\Gamma_\lambda(M \otimes V_{\mu-\lambda})]_\mu = [\widetilde{R}\Gamma(M \otimes V_{\mu-\lambda})]_\mu \simeq \widetilde{R}\Gamma_\mu([M \otimes V_{\mu-\lambda}]_\mu) \quad (4.5)$$

Note also that that $M \otimes V_{\mu-\lambda}$ is an extension of terms $(M \otimes k_{-\nu}) \otimes V_{\mu-\lambda}(\nu)$ (see [BK1] last section) and hence $[M \otimes V_{\mu-\lambda}]_\mu$ is an extension of terms $(M \otimes k_{-\nu}) \otimes V_{\mu-\lambda}(\nu)$ with $\nu \in \mathcal{W}_\lambda^\mu - \lambda \subseteq \mathcal{W}_{\lambda-\mu}$.

If λ, μ lie in the same closed alcove then

$$\mathcal{W}_\lambda^\mu = (\lambda + \mathcal{W}_{\mu-\lambda}) \cap W_{aff} \bullet \mu = (W_{aff})_\lambda \bullet \mu \subseteq \lambda + W \cdot (\mu - \lambda) \quad (4.6)$$

and if μ is also in the closure of the facet of λ then since $(W_{aff})_\mu \subseteq (W_{aff})_\lambda$ we get that $\mathcal{W}_\lambda^\mu = \{\mu\}$. Hence we get from the previous discussion:

Lemma 4.9. *Under the above conditions on μ, λ we have*

$$T_\lambda^\mu(\widetilde{R}\Gamma_\lambda(M)) \simeq \widetilde{R}\Gamma_\mu(M \otimes k_{\lambda-\mu}) \quad (4.7)$$

5 Dimensions of $U_{q,\chi}$ -modules

In this section we follow [BMR]. Set $R = \prod_\alpha < \rho, \check{\alpha} >$ where α runs over the set of positive roots.

Theorem 5.1. *Fix χ a nilpotent element in the big cell and a regular $\lambda \in P$. For any module $M \in U_q - \text{mod}^{\lambda,\chi}$ there exists a polynomial $\mathbf{d}_M \in \frac{1}{R}\mathbb{Z}[P^*]$ of degree $\leq \dim(\mathcal{B}_\chi)$, such that for any $\mu \in P$ in the closure of the alcove of λ , we have*

$$\dim(T_\lambda^\mu(M)) = \mathbf{d}_M(\mu) \quad (5.1)$$

$\mathbf{d}_M(\mu) = l^{\dim(\mathcal{B})} \mathbf{d}_M^0(\frac{\mu+\rho}{l})$ for another polynomial $\mathbf{d}_M^0 \in \frac{1}{R}\mathbb{Z}[P^*]$, such that $\mathbf{d}_M^0(\mu) \in \mathbb{Z}$ for $\mu \in P$.

Remark 5.2. Without loss of generality we can assume that $\chi \in \mathfrak{n}$ (Using the infinite dimensional group of automorphisms of [CKP]).

Corollary 5.3. *The dimension of any $M \in U_q\text{-mod}^\chi$ is divisible by $l^{\text{codim} \mathcal{B}_\chi}$*

proof We can choose a regular λ and μ in the closure of the λ -facet such that $N = T_\lambda^\mu(M)$. Then by theorem 5.1 $\dim(N) = l^{\dim \mathcal{B}} \mathbf{d}_M^0(\frac{\mu+\rho}{l})$. But then since R is prime to p for $p > h$ we get that the denominator of $\mathbf{d}_M^0(\frac{\mu+\rho}{l})$ is a power of p less than $\deg(\mathbf{d}_M^0) \leq \dim(\mathcal{B}_\chi)$ and hence $\dim(N)$ is divisible by a power of p greater than $\dim(\mathcal{B}) - \deg(\mathbf{d}_M^0) \geq \text{codim}(\mathcal{B}_\chi)$.

Lemma 5.4. *Let $\mathcal{M}_\chi^\lambda$ be the splitting bundle for the restriction of the Azumaya algebra $\mathcal{D}_q^\lambda$ to $\mathcal{B}_\chi$. We have an equality in $K^0(\mathcal{B}_\chi)$:*

$$[\mathcal{M}_\chi^\lambda] = [Fr_* \mathcal{O}_q(l\rho + \lambda)|_{\mathcal{B}_\chi}] \quad (5.2)$$

proof As $\mathcal{M}_\chi^\lambda = \mathcal{M}_\chi^0(\lambda)$ by the construction of the splitting it is enough to check for $\lambda = -\rho$. On the zero section (that is on $\mathcal{B}_0$) the Azumaya algebra is split with splitting bundle $\mathcal{O}_\mathcal{B}^{l\dim \mathcal{B}} \simeq Fr_*(\mathcal{O}_q((l-1)\rho))$. We also have that $U_q^{-\rho}|_{\mathfrak{n}} \simeq \text{End}(M_{-\rho})$, hence if we denote by $\tilde{\mathfrak{n}}$ the preimage of $\mathfrak{n}$ under the Springer map then there exists a splitting bundle $\tilde{\mathcal{M}}$ whose restriction to $\mathcal{B}_\chi$ is $\mathcal{M}_\chi^{-\rho}$. Now we can use the map $\mathcal{B}_\chi \times \mathbb{A}^1 \rightarrow \tilde{\mathfrak{n}}$ given by $(x, t) \mapsto (x, t\chi)$ to pull back this splitting bundle $\tilde{\mathcal{M}}$. By the rational invariance of K^0 we get the result. $\square$

Lemma 5.5. *For any $\mathcal{F} \in D^b(\text{Coh}(\mathcal{B}))$ or $\mathcal{F} \in D^b(\text{Coh}(\mathcal{B}_q))$ there exists a polynomial $\mathbf{d}_\mathcal{F} \in \frac{1}{R}\mathbb{Z}[P^*]$ such that for $\lambda \in P$ the Euler characteristic of $R\Gamma(\mathcal{F} \otimes \mathcal{O}(\lambda))$ equals $\mathbf{d}(\lambda)$. We have in the case of $\mathcal{B}$*

$$\deg(\mathbf{d}_\mathcal{F}) \leq \dim(\text{supp}(\mathcal{F})). \quad (5.3)$$

If $\mathcal{F} \in D^b(\text{Coh}(\mathcal{B}))$

$$\mathbf{d}_{Fr^*(\mathcal{F})}(\mu) = l^{\dim \mathcal{B}} \mathbf{d}_\mathcal{F}\left(\frac{\mu + (1-l)\rho}{l}\right) \quad (5.4)$$

proof The first statement is true for line bundles by the Weyl character formula (and its quantum version) and the fact that the K -group is generated by classes of line bundles (also in the quantum case). The inequality follows from Grothendieck-Riemann-Roch ([BMR]). To prove 5.4 it is enough to check it for $\mu = l\nu - \rho$. Then

$$Fr_*(Fr^*(\mathcal{F})(l\nu - \rho)) \simeq Fr_*(Fr^*(\mathcal{F}(\nu))(-\rho)) \simeq \mathcal{F}(\nu) \otimes Fr_*(\mathcal{O}(-\rho)) \quad (5.5)$$

but from 3.7 we get

$$\mathcal{F}(\nu) \otimes Fr_*(\mathcal{O}(-\rho)) \simeq \mathcal{F}(\nu - \rho)^{\oplus l^{\dim(\mathcal{B})}}. \quad (5.6)$$

$\square$

5.1 proof of the theorem

Let $\mathcal{F}_M \in Coh(\tilde{\mathcal{B}}_\chi)$ be such that $\mathcal{L}^\lambda M \simeq \mathcal{F}_M \otimes \mathcal{M}_\lambda$. The existence of such follows from the localization theorem and the Azumaya splitting over the fibers. Let $[\mathcal{F}_M] \in K(Coh(\tilde{\mathcal{B}}_\chi)) = K(\mathcal{B}_\chi)$ be its class. By Lemma 4.9 we have

$$T_\lambda^\mu(M) = R\Gamma(\mathcal{L}^\lambda(M)(\mu - \lambda)) = R\Gamma(\mathcal{F}_M \otimes \mathcal{M}_\lambda(\mu - \lambda)) = R\Gamma(\mathcal{F}_M \otimes \mathcal{M}_\mu) \quad (5.7)$$

Denote by $\int$ the Euler characteristic of $R\Gamma$. We have

$$\dim(T_\lambda^\mu(M)) = \int_{\mathcal{B}_\chi} [\mathcal{F}_M] \cdot [\mathcal{M}_\mu] \quad (5.8)$$

where the multiplication is the action of K^0 on K . By Lemma 5.4 we have

$$\int_{\mathcal{B}_\chi} [\mathcal{F}_M] \cdot i^*[Fr_*\mathcal{O}_{q,l\rho+\mu}] = \int_{\mathcal{B}} i_*[\mathcal{F}_M] \cdot [Fr_*\mathcal{O}_{q,l\rho+\mu}] = \int_{\mathcal{B}_q} Fr^*(i_*[\mathcal{F}_M(l\rho + \mu)]) \quad (5.9)$$

where $i : \mathcal{B}_\chi \hookrightarrow \mathcal{B}$ is the inclusion. By Lemma 5.5 we get

$$\dim(T_\lambda^\mu) = \mathbf{d}_{Fr^*(i_*\mathcal{F}_M)}(l\rho + \mu) = l^{\dim \mathcal{B}} \mathbf{d}_{i_*\mathcal{F}_M}\left(\frac{\mu + \rho}{l}\right). \quad (5.10)$$

So by 5.3 and 5.4 the polynomial $\mathbf{d}_{i_*\mathcal{F}_M}$ has the required properties.

References

- [ABG] S. Arkhipov, R. Bezrukavnikov, V. Ginzburg, *Quantum groups, the loop Grassmannian, and the Springer resolution.*, J. Amer. Math. Soc. 17 (2004), 595-678.
- [BK1] E. Backelin and K. Kremnizer *Quantum flag varieties, equivariant quantum $\mathcal{D}$ -modules, and localization of Quantum groups.*, Adv. in Math. 203 (2006) 408-429.
- [BK2] E. Backelin and K. Kremnizer *Localization for quantum groups at a root of unity.*, math.RT/0407048.
- [BMR] A. Bezrukavnikov, I. Mirkovic and D. Rumynin, *Localization for a semi-simple Lie algebra in prime characteristic.*, arXiv:math.RT/0205144.

- [CKP] C. De Concini, V.G. Kac and C. Procesi, *Quantum coadjoint action*, JAMS v.5, number 1, (1992) 151-189.
- [CP] C. De Concini, C. Procesi, *Quantum groups*, LNM 1565, 31-140.
- [JL] A. Joseph, G. Letzter *Local finiteness for the adjoint action for quantized enveloping algebras*, J. Algebra 153, (1992), 289-318.